Main Panel

Max Pro Soybean

Reg. No./Nr. L11687, Act No./Wet Nr. 36 of/van 1947
READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Bacterial legume inoculant for effective nitrogen fixation on soybean (*glycine max*) species.	**Bakteriese peulgewas inokulant vir effektiewe stikstoffiksasie op sojaboon (*glycine max*) spesies.**

ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:

Live legume bacteria for soybeans (*Bradyrhizobium japonicum*)
Minimum 6.5 x 10^9 cfu/mℓ

NET VOLUME…….**200 mℓ, 400 mℓ, 1 ℓ, 2 ℓ, 4 ℓ, 6 ℓ, 8 ℓ, 10 ℓ, 12 ℓ**..... **NETTO VOLUME**

NOT CLASSIFIED AS HAZARDOUS IN TERMS OF GHS

GHS Precautionary Statements
Wear protective gloves, protective clothing, eye protection and face protection.
Store at temperatures not exceeding 25°C.

REGISTRATION HOLDER / REGISTRASIEHOUER:

AECI® Limited /AECI® Beperk
Reg. No. / Nr. 1924/002590/06
P.O. Box / Posbus 14418
Witfield, 1467
Tel : (011) 823 8000/ 021 858 1140
www.aeciworld.com/www.aeciph.com

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL: 0861 555 777/082 4468946
Spill Tech: 086 100 0366 (a/h)

Expiry Date / Vervaldatum:

Batch No: **Lotnommer:**

UN No. / VN Nr.: NOT REGULATED / NIE GEREGULEER NIE

Pamphlet

Max Pro Soybean

Reg. No. L11687, Act No. 36 of 1947
READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Bacterial legume inoculant for effective nitrogen fixation on soybean (*glycine max*) species.

ACTIVE INGREDIENT:

Live legume bacteria for soybeans (*Bradyrhizobium japonicum*)
Minimum 6.5 x 10^9 cfu/mℓ

REGISTRATION HOLDER / REGISTRASIEHOUER:

AECI® Limited /AECI® Beperk
Reg. No.1924/002590/06
P.O. Box 14418
Witfield, 1467
Tel : (011) 823 8000/ 021 858 1140

WARNINGS:

GHS Hazard Statement	Pictogram and Signal word
Not Classified as Hazardous.	Not Classified as Hazardous.

Max Pro Soybean is exempted from residue restrictions and may be used on its own on any crop.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS, THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL, AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. FURTHERMORE, THE REGISTRATION HOLDER, DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT, OR HARM TO MAN OR ANIMAL, OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED, DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

GHS Precautionary Statements
Keep out of reach of children. Do not get in eyes, on skin, or on clothing. Wear protective gloves, protective clothing, eye protection and face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Store in a well-ventilated place. Protect from sunlight. Store at temperatures not exceeding 25°C.

RELEVANT SUBSTANCES:

Substance name (IUPAC/EC)	CAS –No. EC-No.	Concentration % by weight	Classification EC1272/2008
Bradyrhizobium japonicum	N/A	1-30%	Not classified

NOTE: The other ingredients not included above, do not cause or contribute towards the correct GHS classification of **Max Pro Soybean** and are therefore, in terms of the South African Regulations for Hazardous Chemical Agents - 2021; Regulation 14(b), not listed.

FIRST AID TREATMENT:

In case of inhalation: If inhaled, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get medical attention if you feel unwell.

In case of skin contact: In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse. Thoroughly clean shoes before reuse. Get medical attention if irritation occurs.

In case of eye contact: Check for and remove any contact lenses. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Cold water may be used. Get medical attention if irritation occurs.

In case of ingestion Do NOT induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Loosen tight clothing such as a collar, tie, belt or waistband. Get medical attention if you feel unwell or are concerned.

NOTE TO PHYSICIAN:

Treat symptomatically.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

SHAKE BEFORE USE

- Do not use for any other legume except as indicated.
- Use immediately once the sachet has been opened.

Application methods: Seed treatment or In-furrow	Dosage	Remarks
Seed treatment: **Max Pro Soybean**	300 mℓ/100 kg of seed	Apply to the seed and mix thoroughly until it is evenly coated.
In-furrow: **Max Pro Soybean**	300 to 450 mℓ/ha	Mix with water and apply in-furrow directly on the seed.

Note: Please ensure the seed is dry before planting starts. Use talc/graphite to shorten this interval.

Tector: We recommend the use of **Tector** with the **Max Pro Soybean** inoculant which will help to increase bacterial survival on the seed or in the soil under most stress conditions. Premix the **Tector** and the **Max Pro Soybean** inoculant and then add to the seed and mix thoroughly until it is evenly coated. Plant treated seed as soon as manageable.

Important: 20-day pre-treatment of seed

Max Pro Soybean inoculant has been manufactured with OSMO-protection technology which allows planting up to 20 days after treatment. Premix with **Tector** and the **Max Pro Soybean** inoculant and add this to the seed and mix thoroughly until all the seed is evenly coated. **Do not** use any further water or other foreign liquids in the application. Once the seed is dry, bag and store in a cool (4-24ºC), dark and well ventilated store.

Note: The seed quality, size, moisture content and germination percentage prior to treatment will affect the seed treatment load quantity and consistency of the seed after treatment. Please ensure that seed quality has been tested prior to treatment. This product was tested on registered and certified seed; and recommendations are made accordingly to ensure quality in seed treatment performances and planting practices.

High humidity may result in seed becoming sticky after treatment. If this occurs, or if the time between mixing and planting needs to be shortened, follow planter recommendations for the use of talc and graphite to dry seed out. If the application of talc/graphites not mentioned in planter recommendations, add as required or until seed is sufficiently dry.